



e-ISSN: 2456-6632

This content is available online at AESA

Archives of Agriculture and Environmental Science

Journal homepage: www.aesacademy.org



ORIGINAL RESEARCH ARTICLE



Response of okra (*Abelmoschus esculentus* L. Moench) to vermicompost, mycorrhiza and micronutrients mixture

Manisha Dhayal^{1*} , Kavita Aravindkshan¹, Manoj Kumar Meena¹, Ashok Choudhary² and Rajesh Choudhary³

¹Department of Vegetable science, ²Department of floriculture, College of Horticulture and Forestry, Jhalrapatan city, Jhalawar-326023 (Rajasthan), INDIA, ³Department of Horticulture, Banaras Hindu University, Varanasi - (Uttar Pradesh), INDIA
*Corresponding author's E-mail: msdhayal90@gmail.com

ARTICLE HISTORY

Received: 23 March 2018
Revised received: 13 April 2018
Accepted: 05 May 2018

Keywords

Arka Anamika
Micronutrients mixture
Mycorrhiza
Okra (*Abelmoschus esculentus*)
Vermicompost

ABSTRACT

A field experiments on okra (*Abelmoschus esculentus* L Moench) cv. Arka Anamika" was conducted at Department of Vegetable Science, College of Horticulture and Forestry, Jhalrapatan city, Jhalawar during the Kharif 2016-17 to study the effect of vermicompost, mycorrhiza and micronutrients mixture of 16 treatment combination i.e. vermicompost with two levels (2 tons/ha and 4 tons /ha), mycorrhiza with one level (Mycorrhiza seed treatment) and micronutrients mixture with two levels (50 ppm and 100 ppm). The observations revealed that the plant height of *A. esculentus* after 60 DAS (74 cm), plant height after 90 DAS (79.71cm), number of leaves / plant (43.17), main stem diameter (5.52 cm), number of primary branches per plant (5.10) of *A. esculentus* and lowest days to first flower bud emergence (33.53) of *A. esculentus* was noted maximum as compared to control. The results indicates that plant height, number of leaves per plant, main stem diameter, stem girth, number of primary branches increased significantly due to application of different levels of vermicompost, mycorrhiza and micronutrients mixture as compared to control. It is postulated that the vermicompost 2tons and 4tons + mycorrhiza seed treatment + micronutrients mixture 50ppm and 100ppm may positively regulate the *A. esculentus* growth improved. Therefore, application of vermicompost, mycorrhiza along with a mixture of micronutrients played a significant role on enhancing the growth of okra (*A. esculentus*) and can be applied to obtain the maximum crop yield of okra (*A. esculentus*).

©2018 Agriculture and Environmental Science Academy

Citation of this article: Dhayal, M., Aravindkshan, K., Meena, M.K., Choudhary, A. and Choudhary, R. (2018). Response of okra (*Abelmoschus esculentus* L. Moench) to vermicompost, mycorrhiza and micronutrients mixture. *Archives of Agriculture and Environmental Science*, 3(2): 198-201, https://dx.doi.org/10.26832/24566632.2018.0302014

INTRODUCTION

Okra (*Abelmoschus esculentus* L. Moench) popularly known as 'Bhendi' is an important warm season vegetable, widely cultivated for its tender, green fruits. There are two cultivated types of okra (*Abelmoschus esculentus* L. Moench) and West African okra (*Abelmoschus caillei*). In India, okra mostly grown during Kharif (Chopra *et al.*, 2013; Kumar and Chopra, 2013). It is also mature during premature winter period of mid cool temperatures when cost leftovers very high and farmers earn handsome remuneration from such an okra crop. Okra plays an important role in the diet by supplying carbohydrates, protein, fat, minerals and

vitamins that are usually deficient in the staple food (Kumar *et al.*, 2017). Okra is basically low in calories and dry matter constituents. Generally, it is consumed in a meal with basic starchy food makes the food more palatable (Savello *et al.*, 1982; Ilodibia *et al.*, 2017). Vermicompost acts both as plant growth promoter and plant growth protector. It keeps plants against numerous pests and diseases either by suppressing or resisting them or by inducing biological resistance in plants to fight them or by killing them through pesticide action (Sinha *et al.*, 2010). Mycorrhiza is the edifices resulting from the association between fungi and plant roots and directly involved in plant mineral nutrition. The interdependent root-fungal association

increases the uptake of less movable nutrients (Ortas *et al.*, 2001). Zinc, Boron and Copper are essential trace element for plants being necessary for their good growth and development and involved in many enzymatic reactions. The role of Zinc in plant is due to its requirement in the synthesis of tryptophan (amino acid) which is a precursor of indole acetic acid (IAA) and formation of this growth substance is directly influenced by Zinc. Copper (Cu) is required for physiological redox processes, pollen viability and lignification (Marschner, 1995; Chopra *et al.*, 2017). Therefore, the present investigation was carried out to study the response of okra (*Abelmoschus esculentus* L. Moench) to vermicompost, mycorrhiza and micronutrients mixture.

MATERIALS METHODS

Experimental design and cultivation practices

A field experiment was conducted at the Department of Vegetable Science, College of Horticulture and Forestry, Jhalrapatan city, Jhalawar during Kharif 2016-17. The soil organic carbon 0.56%, available nitrogen 336.29kg/ha, available phosphorous 22.72kg/ha, available potash 220.69kg/ha. The direct seed sowing of cv. Arka Anamika at the spacing 45 × 45 cm by check-basin method in drip irrigation in randomized block design (RBD). A total 15 treatments (T₁- T₁₅) along with control were used for the cultivation of okra (*Abelmoschus esculentus* L. Moench). The description of different treatments is given in Table 1. The observations recorded on number of the plant height after 60 days after sowing (DAS), plant height after 90 DAS, number of leaves / plant, main steam diameter, primary branches per plant, days to first flower bud emergence.

Statistical analysis

The experimental data was expressed as mean ± S.E. One way analysis of variance (ANOVA) and least significant difference (RBD) was carried out using MS Excel to determine difference from control and between the treatments ($P \leq 0.05$).

RESULTS AND DISCUSSION

The effects of vermicompost, mycorrhiza and micronutrients mixture on various growth parameters viz., plant height, number

of leaves, stem diameter, number of primary branches and flower bud emergence of okra (*A. esculentus*) are presented in Table 2.

The result of present study clearly indicates that plant height, number of leaves per plant, main stem diameter, stem girth, number of primary branches increased significantly due to application of different levels of vermicompost, mycorrhiza and micronutrients mixture as compared to control. The maximum value of growth parameters i.e. plant height both at 60 and 90 DAS (74.00 and 79.71 cm), number of leaves per plant (43.17), main stem diameter (5.52 cm) number of primary branches (5.10). and Lowest day to first flower bud emergence (33.53) of okra (*A. esculentus*) was recorded under treatment T₁₅ (vermicompost 4 tons/ha + mycorrhiza + micronutrients mixture @ 100 ppm) and the minimum value of growth parameters i.e. plant height both at 60 and 90 DAS (64.20 and 67.66 cm), number of leaves per plant (34.53), main stem diameter (3.50 cm), number of primary branches (2.77) of okra (*A. esculentus*) was minimum, and highest day to first flower bud emergence (40.73) were recorded under control (T₀). These findings clearly indicated that combination of vermicompost, mycorrhiza and micronutrients mixture played a significant role on enhancing the growth of okra (*A. esculentus*). These results are in close proximity with earlier researches of Mohandas (1987) in tomato, Vadiraj *et al.* (1998) in turmeric, Singh *et al.* (2002) in tomato, El-Rohidas *et al.* (2010) in garlic, El-Tohamy *et al.* (2009) in onion, Ramakrishnan and Selvakumar (2012) in tomato, Chowdhury *et al.* (2014) in okra, and Mehraj *et al.* (2015) in okra. Adequate supply of vermicompost early in the crop season resulted in greater availability of nutrients particular in crop root zone. Increased obtainability of nutrients in the root zone coupled with increased metabolic activity at the cellular level might have increased the nutrient uptake and accumulation in the vegetative plant parts which in turn resulted in improved plant growth attributes (Atiyeh *et al.*, 1999). With vermicompost application of mycorrhiza (bio fertilizer) might have increase higher absorption of nutrients, enhanced cell division, cell elongation and thus concomitant increase in metabolic activity (Torrey, 1950). Micronutrients (Zn) play an important role in many physiological process and cellular formation within the plants. The synthesis of tryptophan in the presence of zinc, the precursor of IAA, which stimulated the growth of plant tissues (Basavarajeshwari *et al.*, 2008).

Table 1. Description of different treatments used for cultivation of okra (*A.esculentus*).

S.N.	Symbols	Treatments
1.	T ₀	Control
2.	T ₁	V ₁ (Vermicompost 2ton /ha)
3.	T ₂	V ₂ (Vermicompost 4ton /ha)
4.	T ₃	M (Mycorrhiza seed treatment)
5.	T ₄	Z ₁ (Micronutrients mixture @50 ppm)
6.	T ₅	Z ₂ (Micronutrients mixture @100 ppm)
7.	T ₆	V ₁ +M (Vermicompost 2ton /ha + Mycorrhiza seed treatment)
8.	T ₇	V ₁ +Z ₁ (Vermicompost 2ton /ha + Micronutrients mixture @50 ppm)
9.	T ₈	V ₁ +Z ₂ (Vermicompost 2ton /ha + Micronutrients mixture @100 ppm)
10.	T ₉	V ₁ +M+Z ₁ (Vermicompost 2ton /ha+ Mycorrhiza seed treatment+ Micronutrients mixture @50 ppm)
11.	T ₁₀	V ₁ +M+Z ₂ (Vermicompost 2ton /ha+ Mycorrhiza seed treatment+ Micronutrients mixture @100 ppm)
12.	T ₁₁	V ₂ +M (Vermicompost 4ton /ha+ Mycorrhiza seed treatment)
13.	T ₁₂	V ₂ +Z ₁ (Vermicompost 4ton /ha+ Micronutrients mixture @50 ppm)
14.	T ₁₃	V ₂ +Z ₂ (Vermicompost 4ton /ha+ Micronutrients mixture @100 ppm)
15.	T ₁₄	V ₂ +M+Z ₁ (Vermicompost 4ton /ha+ Mycorrhiza seed treatment+ Micronutrients mixture @50 ppm)
16.	T ₁₅	V ₂ +M+Z ₂ (Vermicompost 4ton /ha+ Mycorrhiza seed treatment+ Micronutrients mixture @100 ppm)

Table 2. Effect of vermicompost, mycorrhiza and micronutrients mixture on growth of okra (*A. esculentus*).

Treatment	Plant height 60 DAS (cm)	Plant height 90DAS (cm)	Number of leaves/plant	Main stem diameter (cm)	No. of primary branch/plant	Days to first flower bud emergence
T ₀ Control	64.20	67.66	34.53	3.50	2.77	40.73
T ₁ (V ₁)	65.93	68.72	37.77	3.51	3.07	39.80
T ₂ (V ₂)	66.17	68.90	39.21	3.63	3.47	39.30
T ₃ (M)	68.99	71.11	38.45	3.73	3.07	39.10
T ₄ (Z ₁)	68.38	69.85	40.49	3.52	3.10	38.60
T ₅ (Z ₂)	68.62	70.82	35.51	3.71	3.40	38.00
T ₆ (V ₁ +M)	69.08	73.73	40.09	4.02	3.63	37.63
T ₇ (V ₁ +Z ₁)	67.68	71.96	40.44	4.00	3.53	37.50
T ₈ (V ₁ +Z ₂)	67.95	72.42	38.90	4.19	3.73	36.48
T ₉ (V ₁ +M+Z ₁)	68.22	74.53	36.62	3.87	3.67	36.54
T ₁₀ (V ₁ +M+Z ₂)	68.95	75.33	38.01	4.37	3.77	36.47
T ₁₁ (V ₂ +M)	70.04	76.66	36.07	4.51	3.97	35.80
T ₁₂ (V ₂ +Z ₁)	70.52	77.77	38.17	4.67	4.13	36.00
T ₁₃ (V ₂ +Z ₂)	71.51	77.93	37.78	4.78	4.47	35.13
T ₁₄ (V ₂ +M+Z ₁)	73.96	78.12	41.79	5.32	4.87	34.23
T ₁₅ (V ₂ +M+Z ₂)	74.00	79.71	43.17	5.52	5.10	33.53
CD at 5%	4.04	4.76	1.01	0.51	0.30	0.74
SEm±	1.98	2.33	0.49	0.25	0.14	0.36

Conclusion

It is postulated that the vermicompost 2tons and 4tons +mycorrhiza seed treatment + micronutrients mixture 50ppm and 100ppm may positively regulate the okra growth improved. The increasing plant height after 60 DAS(74cm), plant height after 90 DAS (79.71cm), number of leaves / plant (43.17), main steam diameter (5.52cm), primary branches per plant (5.10), days to first flower bud emergence (33.53 DAS) okra (*A. esculentus*) over the control. It was clearly evidenced that plant height, number of leaves per plant, main stem diameter, stem girth, number of primary branches increased significantly due to application of different levels of vermicompost, mycorrhiza and micronutrients mixture as compared to control. Therefore, the findings indicated that combination of vermicompost, mycorrhiza and micronutrients mixture played a significant role on enhancing the growth of okra (*A. esculentus*).

Open Access: This is open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

REFERENCES

Atiyeh, R.M., Subler, S., Edwards, S.A. and Metzger, J. (1999). Growth of tomato plant in horticulture potting media amended with vermicompost, *Pedobiologic*, 43 (8):724 -728.

- Basavarajeshwari, Patil.C., Hosamani, R.M.P., Ajjappalavara, P.S., Naik, B.H., Smitha, R.P. and Ukkund, K.C. (2008). Effect of foliar application of micronutrients on growth and yield components of tomato (*Lycopersicon esculentum* Mill.). *Karnataka Journal of Agriculture Science*, 21(3): 428-430.
- Chopra, A.K., Srivastava, S., Kumar, V. and Pathak, C. (2013). Agro-potentiality of distillery effluent on soil and agronomical characteristics of *Abelmoschus esculentus* L. (Okra). *Environmental Monitoring and Assessment*, 185: 6635-6644.
- Chopra, A.K., Payum, T., Srivastava, S. and Kumar, V. (2017). Effects of integrated nutrient management on agronomical attributes of tomato (*Lycopersicon esculentum* L.) under field conditions. *Archives of Agriculture and Environmental Science*, 2(2): 86-91
- Chowdhury, M.S., Hasan, Z., Kabir, K., Jahan, M.S. and Kabir M. H. (2014). Response of Okra (*Abelmoschus esculentus* L.) to growth regulators and organic manures. *The Agriculturist*, 12(2): 56-63.
- El-Tohamy, W.A., Khalid, A.K., El-Abagy, H.M. and Abou-Hussein, S.D. (2009). Essential oil, growth and yield of onion (*Allium cepa* L.) in response to foliar application of some micronutrients. *Australian Journal of Basic and Applied Sciences*, 3(1): 201-205.
- El-Rohidas, S.B., Bharadia, P.S., Jature, S.D. and Ghate, K.B. (2010). Effect of micronutrients on growth and yield of garlic (*Allium sativum* L.) var. G-41. *Asian Journal of Horticulture*, 5(2): 517-519.
- Ilodibia, C.V., Achebe, U.A. and Chiafor, C. (2017). Nutrient characteristics assessment of two variants of okra (*Abelmoschus*

- esculentus* L. Moench.) found in Anambra State, Nigeria. *Archives of Agriculture and Environmental Science*, 2(4): 298-300, <https://dx.doi.org/10.26832/24566632.2017.020408>
- Kumar, V. and Chopra, A.K. (2013). Ferti-irrigational effect of paper mill effluent on agronomical characteristics of *Abelmoschus esculentus* L. (Okra). *Pakistan Journal of Biological Sciences*, 16 (22): 1426-1437.
- Kumar, V., Chopra, A.K., Srivastava, S., Singh, J. and Thakur, R.K. (2017). Irrigating okra with secondary treated municipal wastewater: Observations regarding plant growth and soil characteristics. *International Journal of Phytoremediation*, 19 (5): 490-499, [https://dx.doi.org/ 10. 1080/15226514.2016.1244169](https://dx.doi.org/10.1080/15226514.2016.1244169)
- Marschner, H. (1995). Functions of mineral nutrients: macronutrients, In: H. Marschner. Mineral nutrition of higher plants 2nd Edition. Academic Press, N.Y. London pp. 299-312.
- Mehraj, H., Taufique,., Mandal, M. S. H., Sikder, R.K. and Uddin, A.F.M. (2015). Foliar feeding of micronutrient mixtures on growth and yield of okra (*Abelmoschus esculentus*). *American -Eurasian Journal of Agriculture and Environmental Science*, 15 (11): 2124-2129.
- Mohandas, S. (1987). Field response of tomato (*Lycopersicon esculentum*) cv. Pusa Ruby. *Plant and Soil*, 98: 295-297.
- Ortas, I., Kaya, Z. and Çakmak, I. (2001). Influence of mycorrhiza inoculation on growth of maize and green pepper plants in phosphorus and zinc deficient soils. In: Plant nutrition- Food security and sustainability of agro-ecosystems (Horst W.J. et al., eds). *Kluwer Academic Publication, Dordrecht*. pp. 632-633.
- Ramakrishnan, K. and Selvakumar, G. (2012). Influence of AM fungi on plant growth and nutrient content of tomato (*Lycopersicon esculentum* Mill.). *International Journal of Research in Botany*, 2(4):24-26.
- Savello, P.A., Martin, F.W. and Hull, J.M. (1982). Nutrition composition of okra seed meal. *Journal of Agriculture Food Chemistry*, 28: 1163-1166.
- Singh, N.P. and Joshi, P. (2002). Nutritive and medicinal value of cucurbits. A book of vegetable on cucurbits by ICAR, pp.7-8.
- Sinha, R.K., Agarwal, S., Chauhan, K. and Valani, D. (2010). The wonders of earthworms & its vermicompost in farm production: Charles Darwin's 'friends of farmers', with potential to replace destructive chemical fertilizers from agriculture. *Agricultural Sciences*, 1(2): 76-94.
- Torrey, J.C. (1950). The induction of lateral shoots by IAA and decapitation. *American Journal of Botany*, 37: 2557-2564.
- Vadiraj, B.A. and Siddagangaiah, Potty, S.N. (1998). Response of coriander (*Coriandrum sativum* L.) cultivars to garden level of vermicompost. *Journal of Spices and Aromatic Crop*, 7: 141-143.

The growth response and nutrient concentration in okra (*Abelmoschus esculentus* L. Moench) as influenced by four nitrogen rates (0, 25, 50 and 100 kg ha⁻¹) and three phosphorus rates (0, 13 and 26 kg ha⁻¹) were examined using two varieties (‘White velvet’ and ‘NHAE 47-4’). Nitrogen application generally increased fruit and shoot dry weights markedly whereas phosphorus increased them only moderately. Leaf and primary branch production and plant height were also enhanced by nitrogen fertilization up to 100 kg N ha⁻¹ but were not influenced by phosphorus application. The application of nitrogen in Okra *Abelmoschus esculentus* L. (Moench) is one of the most widely cultivated vegetables by rural farmers of Nigeria due to its high nutritional value. The potentials of biofertiliser as an alternative to inorganic fertilisers in crop production has been identified. Thirteen treatments combinations of lime, NPK doses and vermicompost were imposed in pea-okra cropping system from rabi 2008-09 to kharif 2011. The recommended doses in 100 % N, P and K were 50:11:42 and 75:21:42 for pea and okra, respectively. The results revealed that addition of 125% [Show full abstract] NPK with 10 t vermicompost ha⁻¹ recorded highest pod/fruit yield of pea and okra during all the years as compared to rest of the treatments. Growth response of okra (*abelmoschus esculentus* (L) moench) to arbuscular mycorrhizal fungus inoculation in sterile and non-sterile soil. Journal: IMPACT : International Journal of Research in Applied, Natural and Social Sciences (IMPACT : IJRANSS) (Vol.2, No. 11). Publication Date: 2015-01-03. Authors : INIOBONG E. OKON; Page : 183-190. Keywords : Okra; Mycorrhiza; Sterile and Non-Sterile Soil; Source : Download Find it from : Google Scholar. Abstract. In a pot experiment, the growth of okra inoculated with *Glomus mosseae* in sterile and non-sterile soil was investigated. Inoculation with *G.*