

## CLINICAL CASE

---

### Epidural metastasis of Ewing's sarcoma: a rare localisation

F.O. Dincbas<sup>1</sup>, D.C. Oksuz<sup>1</sup>, B. Atalar<sup>1</sup>, S. Dervisoğlu<sup>2</sup>, N. Kocer<sup>3</sup>, N.M. Mandel<sup>4</sup>, S. Koca<sup>1</sup>

<sup>1</sup>Department of Radiation Oncology, <sup>2</sup>Department of Pathology, <sup>3</sup>Department of Radiology, <sup>4</sup>Department of Medical Oncology, Istanbul University, Cerrahpasa Medical Faculty, Istanbul, Turkey

#### Summary

*Epidural spinal metastasis of Ewing's sarcoma is rarely observed. We report on a rare case of purely epidural spinal metastasis of Ewing's sarcoma with pain and*

*paraplegia, and describe the treatment and final outcome of the patient.*

**Key words:** epidural, Ewing's sarcoma, metastasis, radiotherapy

#### Introduction

Ewing's sarcoma of the bone is the most common malignant bone tumor in children younger than 10 years of age [1,2]. It is a small round-cell tumor of the bone [3,4]. Metastatic spread is frequent, lungs and bones being the most common sites of metastases [5]. Although Ewing's sarcoma metastases can involve almost every site, epidural metastasis is only rarely observed. We report on a case of Ewing's sarcoma metastatic to the epidural space.

#### Case presentation

A 10-year-old boy had been presented with a mass on the first metatarsal of the left foot with a histologic diagnosis of Ewing's sarcoma in 1992. He was treated with 14 courses of chemotherapy (doxorubicin

20 mg/m<sup>2</sup>/day, days 1-3, plus vincristine 1.4 mg/m<sup>2</sup>, day 1, alternating every 3 weeks with actinomycin-D 0.5 mg/m<sup>2</sup>/day, days 21-23, plus ifosfamide 2 g/m<sup>2</sup>/day, days 21-23, with mesna (protocol EICESS 92) [6], and he then received 54.4 Gy local radiotherapy. After 4-year disease-free survival, a solitary metastatic mass in the right lung was diagnosed. He was treated with right middle lobectomy and combination chemotherapy with cyclophosphamide 1800 mg/m<sup>2</sup> plus vincristine 1.4 mg/m<sup>2</sup> every 3 weeks, alternating with ifosfamide 1.8 g/m<sup>2</sup>/day, days 1-3 plus mesna, and etoposide 100 mg/m<sup>2</sup>/day, days 1-5 every 3 weeks for one year. During follow-up, a new metastasis was found in the lower lobe of the right lung in October 2001. Right lower lobe wedge resection was performed and the patient received 6 courses of continuous infusion ifosfamide 2.5 g/m<sup>2</sup>/day with mesna, days 1-5. He was then referred to our clinic in February 2002. Bilateral lung irradiation up to 18 Gy was delivered and a boost dose of 45 Gy to the metastatic tumor volume was administered. For a long time the patient was without disease-related symptoms. In the 14th month after lung irradiation he presented with back pain persisting for 1 month. Physical and neurological examination, thorax computed tomography, bone scan and biochemistry were all normal. However, 4 days later he developed weakness and numbness in both legs. Neurological examination was unremarkable, except loss of power in the lower extremities (3/5) and hypoaesthesia below the cervical (C) 8 level. Magnetic resonance imaging (MRI) of the thoracic

---

Received 05-11-2004; Accepted 28-11-2004

*Author and address for correspondence:*

Fazilet Oner Dincbas, MD  
Istanbul University, Cerrahpasa Medical Faculty  
Department of Radiation Oncology  
34303, Aksaray, Fatih  
Istanbul  
Turkey  
Tel: +90 212 414 30 00  
Fax: +90 212 414 30 97  
E-mail: faziletonerdincbas@hotmail.com


spine revealed an homogenous epidural mass at the thoracic (T) 6- T7 region, relatively hyperintense on T2-weighted images and hypointense on T1-weighted images. The right posterolateral epidural location of the mass compressed the cord to the left anterior portion of the spinal canal. The right foraminal extension of the mass also occupied the periforaminal fatty place (Figures 1 and 2). The MRI study of the brain and lumbar vertebrae was normal. Therefore, a T6-T7 laminectomy and gross total removal of the tumor were performed urgently.



**Figure 1.** Magnetic resonance imaging of the thoracic spine mass (arrow).



**Figure 2.** Magnetic resonance imaging of the thoracic spine mass (arrow).



**Figure 3.** Tumor consisting of undifferentiated primitive cells with small round-cell morphology (H&E  $\times 400$ ).

The curettage material of 2 cc volume taken from the extradural T6-T7 space was evaluated histologically. The tumor consisted of undifferentiated primitive small blue round cells (Figure 3). Some areas showed rosetoid configuration of tumor cells. The case was considered as Ewing's sarcoma metastasis, as the patient had a history of this malignancy. Cytological examination of the cerebrospinal fluid for meningeal involvement revealed no atypical cells but macrophages and lymphocytes.

After spinal decompressive surgery his neurological status partially improved. He could walk and stand up with help. The patient received radiotherapy with 30 Gy in 15 fractions to the T5-T8 region postoperatively. However, his neurological deficit progressed during radiotherapy. Neurological examination revealed rectal and urinary incontinence, hypaesthesia below the level of T6-T7 and positive Romberg sign. The patient was put on dexamethasone 8mg i.v.x3/daily. A new craniospinal MRI was done. The cranial MRI was normal; however, the spinal MRI revealed a pathological signal posterior to the corpus of L1 with an anterior epidural mass. The patient received radiotherapy with 30 Gy in 10 fractions to T12-L2 region. Chemotherapy was planned, but it never started because of the platelet count ( $59,000/\text{mm}^3$ ) and his bad general condition. Four days after the radiotherapy started, bilateral abducent nerve paralysis, difficulty in swallowing and loss of hearing developed. Although the cranial MRI was normal, we considered these findings as a cranial infiltration, while he responded clinically to mannitol and dexamethasone treatment. Whole brain irradiation was started and cyclophosphamide  $250 \text{ mg}/\text{m}^2$  days 1-3 was administered but 7 days after chemotherapy the patient died due to haemoptysis and febrile neutropenia.

## Discussion

Ewing's sarcoma is an undifferentiated blue round-cell tumor of the bone. It originates from neural tissue, specifically from postganglionic, parasympathetic, primordial cells [3,4]. Seventy-five to 80% of patients with Ewing's sarcoma present with localized disease at diagnosis. The most common sites of metastases are the lung, bone, bone marrow, soft tissue, brain and spine [5]. According to reports of the Bristol Bone Tumor Register, the initial metastasis in patients with long bone primaries can occur both in pulmonary and extrapulmonary sites. When other bones are involved, there is a tendency for extrapulmonary sites [7]. When the primary tumor is not in the long bones, vertebral metastasis can occur with higher probability due to the Batson's pathway. Mehda et al. reported central nervous system (CNS) metastasis in 15 out of 27 patients (55.5%) [8]. However, results from the IECS and the National Cancer Institute studies did not confirm this high incidence of CNS involvement [9,10]. In the Trigg's et al. review [10] the CNS involvement was 2.2% and in almost all of these patients CNS involvement was associated with metastasis in other sites. Meningeal or intraparenchymal CNS disease was the initial site of recurrence in 3 out of 10 patients.

Extra-or epidural metastases are relatively frequent. However, spinal cord compression with serious complications from malignant diseases in children is rare. Spread pathways of epidural metastases are direct extension of mediastinal or retroperitoneal masses or nodes through the intervertebral foramina or transcortical extension from metastatic disease in the vertebral corpus or haematogenous, paravertebral and epidural plexus venosus [11]. The presenting symptoms are usually sudden pain exacerbation, which is difficult to control with increasing doses of analgesics and worsens with recumbency, cough and Valsalva maneuver, pain with radiculopathy and Lhermitte sign [11]. The onset symptom of our case was only pain, however his pain progressed and paraplegia occurred in just 4 days. The differences in symptoms between epidural and intradural metastases do not differ too much, but they can be distinguished from each other. Bladder and bowel dysfunction is much more present in intradural-extramedullary metastasis than in epidural metastases. Also, prominent pain which is experienced at an early stage is more severe and crampy and radicular dysfunction is more prominent in intradural-extramedullary metastasis. Intramedullary metastases have a relatively rapid progressive symptomatology. The onset may be acute, with paraplegia in a few days. Early diagnosis and prompt treatment to achieve the

best possible clinical result, and to keep the patient ambulatory and continent is very important in spinal epidural metastases.

In the Klein et al. study, consisting of pediatric spinal epidural metastases, 30 (17.9%) of 168 patients with Ewing's sarcoma developed spinal cord compression from epidural tumor spread [12]. In this study, the children's neurologic function rapidly improved both with chemotherapy and radiotherapy without surgery in small-cell tumors (neuroblastoma, germ-cell, Hodgkin's lymphoma). On the contrary, patients with Ewing's sarcoma showed a significant improvement in survival by adding decompressive surgery to treatment. The mean survival was 4.2 years with decompressive laminectomy. In the study of Bacci et al. there were 2 patients with spinal cord compression due to epidural tumor growth, with no radiological evidence of bone involvement [13].

In conclusion, although pure epidural metastases of Ewing's sarcoma are uncommon, they should be included in the differential diagnosis of an epidural mass lesion. Although local regression is obtained after decompressive surgery and radiotherapy, rapid spread of the disease through the spinal canal and to the cranium, even with subclinical disease, should be considered.

## References

1. Young JL Jr, Miller RW. Incidence of malignant tumors in US children. *J Pediatr* 1975; 86: 254-258.
2. Kozlowski K, Beluffi G, Masel J et al. Primary vertebral tumors in children. Report of 20 cases with brief review of the literature. *Pediatr Radiol* 1984; 14: 129-139.
3. Cavazzana AO, Miser JS, Jefferson J et al. Experimental evidence for a neural origin of Ewing's sarcoma of bone. *Am J Pathol* 1987; 127: 507-518.
4. Link M, Grier H, Donaldson S. Sarcomas of bone. In: Fernbach D, Vietti T (eds): *Clinical Pediatric Oncology*. St. Louis, Mosby-Year Book, 1991, p 545.
5. Edward CH. Pediatric sarcomas of bone. In: Gunderson LL, Tepper JE (eds): *Clinical Radiation Oncology* (1st edn). Churchill Livingstone, New York, Edinburgh, London, Philadelphia, 2000, pp 1066-1072.
6. Paulussen M, Ahrens S, Braun-Munzinger G et al. EICES 92 (European Intergroup Cooperative Ewing's Sarcoma Study); Preliminary results. *Klin Paediatr* 1999; 211:276-283.
7. McIntosh DJ, Price CH, Jeffree GM. Ewing's tumor: a study in behavior and treatment in 47 cases. *J Bone Jt Surg* 1975; 57: 331-340.
8. Mehta Y, Hendrickson FR. CNS involvement in Ewing's sarcoma. *Cancer* 1974; 33:859-862.
9. Marsa GW, Johnson RE. Altered pattern of metastases following treatment of Ewing's sarcoma with radiation and adjuvant chemotherapy. *Cancer* 1971; 27: 1051-1054.
10. Trigg ME, Glaubiger D, Nesbit ME. The frequency of isolated

- CNS involvement in Ewing's sarcoma. *Cancer* 1982; 49: 2404-2409.
11. Debois JM (ed).  $T_xN_xM_1$ : The Anatomy and Clinics of Metastatic Cancer. Boston, Kluwer Academic Publ, 2002, pp 150-153.
  12. Klein SL, Sanford RA, Muhlbauer MS. Pediatric spinal epidural metastases. *J Neurosurg* 1991; 74: 70-75.
  13. Bacci G, Toni A, Avella M et al. Long-term results in 144 localized Ewing's sarcoma patients treated with combined therapy. *Cancer* 1989; 63: 1477-1486.

